

Seizure Emergency Care Plan and Medication Orders for School or Care Settings

PARENT/GUARDIAN to complete, SIGN and DATE Below

Child/patient Name:	Birth date:
Parent/Guardian Contact:	Phone:
Emergency Contact:	Phone:
School:	Grade:

Triggers: tiredness illness temperature Other: _____

Seizure Warning (aura) if any: _____

Has patient ever received rescue medication before? _____ **Date:** _____

Antiseizure Medications Taken at Home	What side effects does this patient experience?

Is patient being treated with a ketogenic diet therapy for epilepsy?

No Yes, family will arrange or discuss plans for school meals and snacks.

I give permission for school personnel to share this information, follow this plan, administer medication and care for my child and, if necessary, contact our care team. I take full responsibility for providing the school with prescribed medication and devices. I approve this Seizure Emergency Care Plan for my child.

Parent/Guardian Signature	Date	Nurse/CCHC Signature	Date
---------------------------	------	----------------------	------

HEALTH CARE TEAM to complete, SIGN and DATE Below.

IF YOU SEE THIS:	DO THIS:
<input type="checkbox"/> Convulsive Generalized Tonic Clonic: These seizures may begin with a warning (aura). The patient will lose consciousness. You may see stiffening of the body or rhythmic jerking movements. Convulsive seizures may last 1-5 minutes. Sleepiness and confusion may occur after the seizure.	FOR CONVULSIVE SEIZURES ONLY: <ol style="list-style-type: none"> 1. Time the seizure and record observations. 2. Keep calm. Provide reassurance. Remove bystanders. 3. Protect head, keep airway clear, turn on side. 4. Do not restrain or place anything in mouth. 5. Call 911 if patient is injured or has difficulty breathing. 6. Call guardian. 7. Stay with patient until recovered from seizure. Administer rescue treatments as marked below.
<input type="checkbox"/> Focal: These seizures often begin with a warning (aura). The patient may be partly alert or unconscious. You may see lip smacking, chewing, eye blinking, or picking at clothes. These seizures usually last 1-2 mins. Sleepiness and confusion may occur after the seizure.	
<input type="checkbox"/> Absence: The patient may have sudden changes in alertness. You may see eye flutter or small twitching. Usually last less than 10 secs. These are not an emergency unless clustering for more than 10 minutes without return to baseline.	
<input type="checkbox"/> Febrile seizures: The patient may not be on a daily antiseizure medication. Seizures may occur at the beginning of an illness and can appear similar to other seizure types. Preventative treatment with antipyretics does not reduce incidence.	
<input type="checkbox"/> OTHER: please describe: _____ _____ _____	
<input type="checkbox"/> Child has a history of psychogenic non epileptic events (if selected please provide separate documentation for clarification, these do not require rescue treatments)	FOR ALL OTHER SEIZURE TYPES (BESIDES CONVULSIVE): <ol style="list-style-type: none"> 1. Time the seizure and record observations. 2. Gently guide patient away from danger. 3. Stay with patient and reassure them until recovered from seizure. 4. Do not treat staring that is stopped by a touch/nudge. 5. Call guardian. Administer rescue treatments as marked below.
RESCUE TREATMENTS:	

"This document and the information it contains was created by Children's Hospital Colorado ("CHCO") to serve as a guideline and reference tool for use by CHCO employees while acting within the scope of their employment with CHCO. The information presented is intended for informational and educational purposes only. It is not intended to take the place of your personal physician's advice and is not intended to diagnose, treat, cure or prevent any disease. The information should not be used in place of a visit, call, consultation or advice of your physician or other health care provider. Copyright © Children's Hospital Colorado 2024 All rights reserved. No part of this document may be reproduced without written consent from the author." Rev. 08/24

Implantable devices:
 does patient have an implantable device? If yes, please describe: _____

If convulsive seizure lasts longer than 5 minutes, or focal seizure >10 minutes administer:

Diastat: rectally _____mg can give second Diastat dose if seizures continue after ____minutes.
 Nayzilam nasally _____mg can give second Nayzilam dose if seizures continue after ____minutes.
 Midazolam nasally _____mg to administer half in each nostril. can give second nasal midazolam dose if seizures continue after ____minutes
 Valtoco nasally _____mg can give second Valtoco dose if seizures continue after ____hours

Multistep seizure rescue plan – Please see attached letter for details.

OR If cluster of ____or more seizures in ____minutes, administer:

Clonazepam _____mg
 Diastat: rectally _____mg
 Nayzilam nasally _____mg can give second Nayzilam dose if seizures continue after ____minutes
 Midazolam nasally _____mg to administer half dose in each nostril
 Valtoco nasally _____mg can give second Valtoco dose if seizures continue after ____hours

Multistep seizure rescue plan – Please see attached letter for details.

[] Call 911 immediately if emergency medication is administered

[] Call 911 if seizure does not stop within ____min of giving medication.

Does patient have photo-sensitive epilepsy (ie. can flashing lights trigger seizures)? ____ (Y/N)

Accommodations: Always take seizure action plan and emergency medication for school activities, sports and field trips. Close adult supervision when swimming or climbing.

HEALTH CARE PROVIDER SIGNATURE

PRINT PROVIDER'S NAME

PHONE/FAX

DATE

"This document and the information it contains was created by Children's Hospital Colorado ("CHCO") to serve as a guideline and reference tool for use by CHCO employees while acting within the scope of their employment with CHCO. The information presented is intended for informational and educational purposes only. It is not intended to take the place of your personal physician's advice and is not intended to diagnose, treat, cure or prevent any disease. The information should not be used in place of a visit, call, consultation or advice of your physician or other health care provider. Copyright © Children's Hospital Colorado 2024 All rights reserved. No part of this document may be reproduced without written consent from the author." Rev. 08/24