

IN CARE OF KIDS



Metered Dose Inhalers (MDI) with Spacer and Mouthpiece



1. Remove caps from the spacer and metered dose inhaler (MDI).
2. Shake the inhaler for 3-5 seconds.
3. If the inhaler is new, hasn't been used in a while, or has been dropped it needs to be primed. Spray 4 times to prime. This does not count as a dose of medication.
4. Connect the inhaler and the spacer.
5. Stand or sit up straight.
6. Breathe out, insert mouthpiece between teeth and make a tight seal with your lips.
7. Press down on the MDI once to spray into the spacer.
8. Breathe in slowly & deeply through your mouth. YOU SHOULD NOT HEAR A WHISTLE.
9. If you hear a whistle you are breathing in too fast.
10. Hold your breath for 10 seconds. Breathe out normally.
11. Wait 1 minute and repeat steps 2-9 for each puff prescribed.
12. Remove the MDI from the spacer. Replace the cap on your MDI when finished.
13. If you are using an inhaled steroid, rinse your mouth or brush your teeth after each use.

Cleaning instructions:

- Clean spacer once a week.
- Take apart the spacer.
- Rinse in warm soapy water.
- Rinse with clean water and air dry.
- Do not put in the dishwasher.
- Clean the small hole in the MDI once a week with a wet Q-tip.
- Replace the MDI when the counter reaches 000. It may continue to spray, but no medicine is coming out.

Metered Dose Inhalers (MDI) with Spacer and Facemask

*Facemask should be used with children under the age of 7.



1. Remove caps from the spacer and metered dose inhaler (MDI).
2. Shake the MDI for 2-3 seconds.
3. If the inhaler is new, hasn't been used in a while, or has been dropped it needs to be primed. Spray 4 times to prime. This does not count as a dose of medication.
4. Connect the MDI to the spacer. Attach the mask to the mouthpiece of the spacer.
5. Stand or sit up straight.
6. Place the mask over the child's nose and mouth to make a tight seal.
7. Have the child breathe out.
8. Press the top of the MDI so that it sprays into the spacer. Only press one time.
9. Have your child breathe in and out for 10 seconds.
10. Wait 1 minute and repeat steps 2-9 for each puff prescribed.
11. Remove the MDI from the spacer. Replace the cap on your MDI when finished.
12. If your child used an inhaled steroid, wash face and rinse mouth or brush teeth after each use.

Cleaning instructions:

- Clean spacer once a week.
- Remove mask and take apart the spacer.
- Rinse in warm soapy water.
- Rinse with clean water and air dry.
- Do not put in the dishwasher.
- Clean the small hole in the MDI once a week with a wet Q-tip.
- Replace the MDI when the counter reaches 000. It may continue to spray but no medicine is coming out.

To watch a video on how to use the MDI inhaler, visit: <http://www.childrenscolorado.org/MDI>

Author: Asthma Education Standardization Committee Approved by Patient Education Committee Valid through 2019
The information presented is intended for educational purposes only. It is not intended to take the place of your personal doctor's advice and is not intended to diagnose, treat, cure or prevent any disease. The information should not be used in place of a visit, call or consultation or advice of your doctor or other health care provider.

childrenscolorado.org | Anschutz Medical Campus 13123 East 16th Avenue, Aurora, CO 80045 | 800-624-6553